

During the first 24 hours following your discharge

If you have any further concerns about your surgery, please telephone the ADSU: 0118 322 7622. Fax: 0118 322 8214

Opening hours are 7.30am to 10.00pm Monday-Friday, 7.30am to 4.00pm Saturday. If you need help/advice outside these hours, please telephone the Royal Berkshire Hospital switchboard on 0118 322 5111 and ask for the on-call surgery doctor. Please note that this should be for emergencies only.

After 24 hours, please seek advice from your GP.

For further information about the Trust, visit our website www.royalberkshire.nhs.uk

Royal Berkshire NHS Foundation Trust
London Road
Reading RG1 5AN
Telephone 0118 322 5111

ADSU_483
Adult Day Surgery Unit, June 2016
Review due: June 2018

Advice following laparoscopic (keyhole) hernia repair

Adult Day Surgery Unit

What is a hernia?

A hernia is a bulging of internal organs or intra-abdominal fat through a gap or defect in the muscle wall.

The operation involves the insertion of small tubes (usually 3) through the abdominal wall through which a camera and surgical instruments can be passed. When you wake up, you will see three or four plasters over the insertion sites. These sites will have been stitched or glued.

After the operation

- Please start gently mobilising straight away, and you can resume your normal activity gradually, as you feel able. It is quite safe to start exercising 2 weeks post-operatively if you are comfortable enough to do this.
- Your wound dressings are shower proof, so you can take a shower, but avoid soaking the dressings/ bathing for 4 days post-operatively. Thereafter, the dressings can be removed, and the skin heal uncovered.
- *You have dissolvable stitches. These may take four to six weeks to dissolve.
- *Please make an appointment with the practice nurse at your GP's surgery, for removal of stitches on _____.

* *Delete as appropriate.*

- You may experience some upper abdominal pain and discomfort during the first 24 hours post-operatively, which is mainly due to the use of carbon dioxide gas to inflate your abdomen, The pain can radiate up into your shoulders. This is completely normal. You may also experience some lower abdominal aching and swelling, which may be due to bruising (haematoma), and this should gradually improve over the first week post-operatively. You should

take regular painkillers (such as Paracetamol or Ibuprofen) to help relieve your pain.

- It is not uncommon to experience difficulties in passing water post-operatively, particularly if you have had hernia repairs on both groins and/or you have already experienced problems with urinating in the past. If you are unable to pass water once you have gone home, you must come back to the hospital for assessment.
- If the pain worsens or becomes unbearable or you have any other problem related to your operation, please contact your GP.
- You should stay off work for _____. You may need to refrain from work for longer than this, depending on the healing process. Please make an appointment with your GP if this needs to be reviewed.
- Avoid heavy lifting for four to six weeks.
- Do not drive for three to five days as the operation may have affected your reflexes. Check with your motor insurance company before driving a vehicle.

If there is anything you do not understand or if you have any questions or concerns, please feel free to discuss them with your nurse.