



How to prevent a pressure sore (also known as pressure ulcers and bed sores)

This leaflet is written for patients by patients and is based on experience and medical information.

Prevention is better than cure

- Have you ever sat or laid still for too long and felt uncomfortable?
- What if you were unwell and unable to stand up or move position?

Something as simple as this could become the start of a pressure sore.

Pressure sores are painful, hard to heal and can be fatal. **Pressure sores can nearly always be prevented.**


What is a pressure sore?

Pressure sores are caused by excess pressure on the skin, which stops blood and oxygen reaching the skin. This causes a wound to develop. Moisture, friction (including rubbing) and shearing (when skin is moved against a fixed surface) increase the risk of developing a pressure sore.

A pressure sore can start as a painful inflamed area and develop into a large deep wound, down to the underlying bone.

Pressure ulcers often occur from sitting or lying too long in one position, but they can also develop under medical devices, such as masks, plaster casts and nasogastric (NG) tubes.

Pressure damage can occur quickly, within a few hours. If you are concerned about an area of your skin, either because a change in colour, pain in the area, or if you notice the skin is blistered or broken, contact your healthcare professional to seek advice. A change in colour may look different on different skin tones.



© National Wound Care Strategy Programme (NWCSP) 2024

What people say

“ Even when aware of the risk, having had a pressure sore before, and knowing my body best, it felt hard to advocate for myself with hospital staff. ”

“ I wish I'd have known about this and could have prevented it. ”

You are at higher risk of pressures sores if you...

- ⇒ Are unable to move some or all of your body (e.g. paralysis, following stroke or spinal cord injury or MS or Parkinson's disease).
- ⇒ Are spending more time in bed or sitting down due to poor health or following illness or surgery.
- ⇒ Have medical conditions that affect your blood supply (such as diabetes, kidney failure and heart failure).
- ⇒ Are over 70.
- ⇒ Are unable to feel pain or discomfort.
- ⇒ Are dehydrated.
- ⇒ Have a poor diet.
- ⇒ Are very overweight or very thin.
- ⇒ Have urinary and/or bowel incontinence.

It can take years for a pressure sore to heal!

Even when a wound heals, the skin is not the same as before.

Where to get further information and support

- ⇒ Speak to your nurse if you have any questions or concerns.
- ⇒ Visit the NHS Website: www.nhs.uk/conditions/pressure-sores
- ⇒ View Government guidance Pressure Ulcers: Applying All our Health: www.gov.uk/government/publications/pressure-ulcers-applying-all-our-health/pressure-ulcers-applying-all-our-health



Please ask if you need this information in another language or format.