



Alcohol pelvic sympathectomy

This leaflet outlines what to expect during alcohol pelvic sympathectomy treatment in the Pain Management Unit at the Royal Berkshire Hospital.

What is an alcohol pelvic sympathectomy?

An alcohol pelvic sympathectomy is an injection of medicinal alcohol and local anaesthetic around the bundle of nerves that supplies sensation to the veins of the pelvis. Those same nerves can also transmit sensation of pain arising in the organs of the pelvis. It is found at the base of the back in front of the spine and behind the main veins to and from the heart. A steroid may be injected to help reduce inflammation caused by the procedure. The steroid we use is not licensed by the Medicines Control Agency for this procedure, but it has been used without any problems in millions of patients worldwide.

Why am I having this procedure?

You have already had a pelvic sympathectomy with local anaesthetic that gave you a degree of relief that you and your doctor thought was significant. Unfortunately, it has now worn off. Alcohol can be much longer lasting than standard injections.

Is this the right treatment for me?

Other treatment options will be discussed with you before deciding to go ahead with the injection(s) and your consent is needed. The decision on whether or not to go ahead with the injection(s) is a shared decision between you and your doctor. Your doctor will be able to provide you with up-to-date information about the likelihood of this being a successful treatment for you and how this treatment fits into the best pathway of care. If you are undecided about whether or not to have injection(s) then further advice and information to make this informed decision can be provided. Please speak to your doctor for more information.

If your health has changed, it is important to let your doctor know:

- If you have an infection in your body or on the skin of your back, your doctor will postpone the treatment until the infection is cleared.
- If you have been started on anticoagulant or antiplatelet medicines that “thin the blood” such as warfarin, heparin, apixaban or clopidogrel, this may require extra preparation.
- If you suffer from diabetes, the use of steroids during injections may cause your blood sugar to change requiring monitoring and adjustment of your diabetic medication.
- If you have any allergies.
- You must also inform the doctor if there is any chance that you could be pregnant.
- Finally, if you are planning to fly or travel abroad within two weeks after the injections, please let your doctor know.

Are there any risks?

The procedure may fail to provide you with the expected relief. Unfortunately, we don't know if this is going to be the case until we have done it.

Your blood pressure will go down and that can make you feel a little lightheaded.

Occasionally, you may develop muscle spasm in the back. If you take painkillers and continue to move your back normally it should settle within 7 to 10 days.

Other risks are rarer and may include infection, bleeding, nerve damage with pain into your legs, urine retention that may need the use of a catheter, perforation of bowel, veins or kidneys, permanent damage to bowel and permanent diarrhoea.

There is also a small risk that the steroid can temporarily lower your immunity and increase your risk of catching Covid-19.

During the procedure it is normal to feel a painful sensation in your groin. This is from the needle going through some muscles and will settle within 12 hours. If there are any untoward complications we may need to admit you into hospital.

What will happen between now and the injection?

The doctor may have prescribed some painkillers for you to take and unless they are giving you side effects or not helping, you should continue to take them as advised.

What will happen on the day of the injection?

After you have been checked in by a nurse, your doctor will explain the procedure to you and offer you the chance to ask questions. Sedation is not normally required for this procedure, but if you feel particularly anxious, it may be offered after discussion with your doctor. Then you will be asked to sign a consent form.

After that you will be escorted into the treatment room. A small plastic needle may be inserted into the back of your hand. Then you will be asked to lie on your front on the procedures table. Your back will be cleaned with antiseptic solution and some local anaesthetic will be injected into your skin. This will sting. You may feel a pushing sensation as the needle is put into the right place. An X-ray machine and X-ray dye will be used during the procedure to help identify the correct position of the needle. You may feel pressure as the solution of alcohol and local anaesthetic is injected. Then the needle will be removed and your back will be cleaned.

You will be then transferred onto a wheelchair and taken back to your ward where you can eat and drink. Your blood pressure and pulse will be checked. If your blood pressure and pulse remain within normal and safe limits you will be allowed to go home, once all discharge criteria are satisfied.

How will I feel afterwards?

Your tummy may feel: Warm / Tingly / Numb / Stiff / No different

You may develop: Lightheadedness / Nausea / Flushing / Headache / Flu like symptoms

Important things to remember

- You will need a responsible adult who will take you home and stay with you overnight.
- If you have any special requirements, please contact the Pain Management Unit at least 48 hours in advance.
- Please bring your regular painkillers with you.
- Bring a dressing gown and slippers with you.
- If you are on any blood thinners then you need to discuss this with your doctor regarding stopping it, before your procedure.

Can I drive afterwards?

You should not drive for 24 hours after the injection, as your insurance may be void if you are in an accident.

When can I shower/bathe?

Usually, the day after the procedure.

When can I return to normal activity?

Although your pain may feel better from the beginning, the alcohol may take up to 48 hours to work. Your back may take up to 2 weeks to recover. You may do gentle physical activity for 24 to 48 hours following the procedure. If your job is physical up to 5 days of gentle activity is advisable. If your job is sedentary, you may return within 48 hours. However, each individual's response may vary following the procedure.

Any queries?

If you have any concerns regarding this treatment please contact:

During working hours: Pain Management Unit 0118 322 8261.

Out of hours / weekends: please contact your GP or the out of hours service for advice.

To find out more about our Trust visit www.royalberkshire.nhs.uk

Please ask if you need this information in another language or format.

RBFT Pain Management Unit, June 2024.

Next review due: June 2026.