

Inserting a percutaneous biliary drain and biliary stent (a tube to drain bile)

Introduction

This leaflet tells you about the procedures known as percutaneous biliary drainage and biliary stent insertion. It explains what is involved and what the risks are. It is not meant to replace informed discussion between you and your doctor, but to act as a starting point for such a discussion.

Whether you are having the percutaneous biliary drainage or stent insertion as a planned or an emergency procedure, you should feel you have received sufficient explanation before you sign the consent form.

What is percutaneous biliary drainage and biliary stent insertion?

One of the normal functions of the liver is to produce bile. This drains through a series of small tubes, or ducts, into one larger tube, the common bile duct, which then empties into the duodenum, the first part of the bowel after the stomach. If the bile duct becomes blocked, then bile cannot drain normally and jaundice develops. This is potentially a very serious condition, which needs to be investigated promptly and is usually treated.

One of the treatments involves inserting a fine plastic drainage tube through the skin, using only a small incision, into the obstructed bile duct to allow the bile to drain externally for a while. This procedure is called percutaneous (meaning through the skin) biliary drainage.

Sometimes, the biliary drainage procedure may be extended by placing a permanent plastic or metal stent across the blockage in the bile duct. Stents are usually inserted a few days after the initial drainage procedure and they keep the narrowed duct open without the need for a catheter.

To help insert the stent, biliary dilatation might be necessary. This involves stretching a segment of bile duct with a balloon to open up the narrowing either before or after inserting the stent.

The procedure(s) will be done under a local anaesthetic and sedation (you are awake but drowsy, and the area is numb).

Why do I need a percutaneous biliary drainage?

Biliary drainage is necessary if other tests, such as an ultrasound or a CT scan, show that the bile duct has become blocked. Common causes of obstruction are gallstones or tumours in or around the bile duct, liver, gallbladder or pancreas, but sometimes the cause is unknown.

The underlying cause for the obstruction may only become evident once the biliary drainage has been carried out.

Are there other options?

In some cases, as an alternative to percutaneous drainage, biliary drainage can be carried out by passing a flexible telescope, or endoscope, into the small intestine (duodenum) and inserting a drainage catheter into the bile duct that way.

The doctors in charge of your case and the radiologist doing the procedure will have discussed the situation, and will advise on the best option. However, you will also have the opportunity for your opinion to be taken into account, and if, after discussion with your doctors, you do not want the procedure carried out, you can decide against it.

Who will perform the procedure and where will it take place?

The procedure will be performed by an interventional radiologist. This is a doctor specializing in minimally invasive treatments using image guidance. Interventional radiologists are trained to use diagnostic imaging equipment, such as x-ray and ultrasound, to guide various instruments during a procedure. They are supported by radiographers (who are highly trained to carry out x-rays and other imaging procedures) and nurses.

The procedure will take place in the Interventional Radiology Suite in the Radiology Department in a room especially adapted with x-ray and ultrasound equipment.

How do I prepare for percutaneous biliary drainage?

You need to be an inpatient in the hospital. You will be asked not to eat for 6 hours before hand, although you may drink water up to 2 hours before and take your usual medications. A cannula will be inserted in a vein in your arm and intravenous antibiotics will be given. You will be asked to put on a hospital gown and will be taken to the Radiology Department.

If you have any allergies, you must let your doctor know. If you have reacted adversely to intravenous contrast medium (the dye used for kidney x-rays and CT scans) in the past, then you must also tell your doctor about this.

What happens during percutaneous biliary drainage & biliary stent insertion?

You will lie on the x-ray table, generally flat on your back. You will have monitoring devices attached to your chest, arm and finger and will receive oxygen through small tubes in your nose. You will be given sedatives and painkillers intravenously through a cannula, usually in your hand.

The interventional radiologist will keep everything sterile and will wear a theatre gown and operating gloves. The skin of your lower chest and abdomen will be cleaned with antiseptic and the rest of your body will be covered with a special fabric called a theatre drape.

The radiologist will use an ultrasound scanner to decide on the most suitable point for inserting the fine plastic tube (the drainage catheter). This is generally between two of your lower ribs, on the right side or in the upper part of your abdomen in the middle. Then your skin and edge of the liver will be numbed with local anaesthetic. When the local anaesthetic is injected, it will sting to start with, but this soon passes off, and the skin and deeper tissues will then be numb.

Later, you may be aware of the needle and then the wire and catheter passing into the liver, and sometimes this is painful. There will be a nurse standing next to you and looking after you. If the procedure does become painful, then they will be able to give you more painkillers through the cannula in your hand or arm. Generally, placing the catheter in the liver only

takes a short time, and once in place it should not hurt at all.

When the radiologist is sure that the needle is in a satisfactory position in one of the bile ducts, a guide wire will be passed through the needle, into the bile duct, which then enables the plastic drainage catheter to be positioned correctly.

The procedure may finish at this stage, with the catheter being fixed to the skin surface, and attached to a drainage bag. Alternatively, it may be possible to advance the wire and catheter through the obstruction, so that the catheter drains the bile internally into the bowel in the normal way.

What is involved in fitting a stent?

In some cases, a permanent metal tube, called a stent, may be placed across the obstruction, to relieve the blockage. A temporary external catheter may be left in place, attached to a drainage bag. This will be removed after a few days when the stent has expanded fully and the bile is draining through the stent into the bowel. Usually the passage that has been made through the liver will be intentionally blocked using small pellets of sponge to reduce the risk of bleeding and bile leakage.

Will it hurt?

Unfortunately, it may hurt a little, for a very short period of time, but any pain you have should be controlled with painkillers during or after the procedure.

How long will the procedure take?

Every patient's situation is different, and it is not always easy to predict how complex or how straightforward the procedure will be. It may be over in 45 minutes, or occasionally it may take longer than 90 minutes. As a guide, expect to be in the x-ray department for about 2-3 hours altogether.

What happens afterwards?

You will be taken back to your ward on a trolley. Nurses on the ward will carry out routine observations, such as taking your pulse and blood pressure, to make sure that there are no problems. You will generally stay in bed for a few hours, until you have recovered.

If you have an external drainage catheter, attached to a bag, it is important that you take care of this. You should not make any sudden movements (for example, getting up out of a chair) without remembering the bag and making sure that it can move freely with you. However, you will be able to lead a normal life with the catheter in place (except submerging in water i.e. bath). The bag needs to be emptied fairly frequently, so that it does not become too heavy. The nurses on the ward will measure the amount of bile draining into the bag as part of your general fluid management.

How long will the catheter stay in and what happens next?

These are questions which only the doctors looking after you can answer. It depends, for example, on whether you have a temporary external drainage catheter in place, or whether a metal stent has been placed across the blockage. You may require further x-rays or scans to check that the obstruction has been relieved, and to try and determine the cause of the obstruction.

Are there any risks or complications?

Percutaneous biliary drainage is a generally a low risk procedure but complications can arise. It may not be possible to perform the procedure. Often this is because, even though the duct is blocked, the bile ducts in the liver may not become dilated and therefore it is difficult to place a needle into an undilated bile duct. If this happens, your doctors will arrange another method of overcoming the blockage.

Sometimes, there is a leak of bile from the bile duct where the tube has been inserted, resulting in a collection of bile inside the abdomen. This can be painful. Generally, once the catheter is draining bile satisfactorily, the leak should stop. However, if this becomes a large collection, it may require an additional percutaneous drainage procedure.

Patients with jaundice are more likely to have difficulties with blood clotting and there may be bleeding from the liver where the catheter is inserted. Rarely, this may require a blood transfusion. On very rare occasions, this may become severe, and require an operation or another radiological procedure to stop it.

Obstructed bile ducts often contain infected bile. The drainage procedure is needed to treat the infection but can itself result in the infection worsening and going into the blood stream (septicaemia). We try to prevent this by giving intravenous antibiotics.

A recent nationwide registry showed that the mortality (risk of dying) whilst in hospital or in the 30 days after a percutaneous biliary drainage or stent insertion was around 16%. This reflects the seriousness of the complications and the poor health of some of the patients having these procedures. Despite this, the procedure is usually safe and results in an improvement in your medical condition.

Finally...

Some of your questions should have been answered by this leaflet, but remember that this is only a starting point for discussion about your treatment with the doctors looking after you.

Please call your GP if you experience any of the following:

- Recurrence of jaundice.
- Temperatures, feeling shivery or other indications of infection.

Other sources of information

British Society of Interventional Radiology www.bsir.org/patients/biliary-drainage-and-stenting/
For further information about the Trust, visit our website www.royalberkshire.nhs.uk

Royal Berkshire NHS Foundation Trust
London Road, Reading RG1 5AN
0118 322 5111 (switchboard)

This document can be made available in other languages and formats upon request.