

Catheter removal following radical prostatectomy

This leaflet is for patients fitted with a urinary catheter following an operation to remove the prostate. It explains what happens when you return to hospital to have the catheter removed and what to expect afterwards. If you have any further questions or worries during this time please contact Hopkins Ward.

On arrival

Before you return to Hopkins Ward for your catheter removal, please make sure you have drunk plenty of fluids. After we remove your catheter please drink as you normally would for your comfort.

What to expect after catheter removal

Catheter removal is quick and painless. Once the catheter has been removed, temporary problems with urinary control are common and it is normal to experience leaking.

However, we will still ask you to try and use a urinal bottle if possible.

Once the catheter has been removed you may experience one or more of the following:

- Frequency of passing urine - It is quite normal to want to pass urine frequently for the first couple of hours after your catheter has been removed. Sometimes, this can be as often as every half hour. This is nothing to be concerned about and should settle over a period of one or two days.
- Urgency - You may get very little warning when you want to pass urine and you may have to hurry to the toilet or you may not make it to the toilet in time. Again this is normal and will settle into a more normal pattern in time.
- Incontinence - It is normal to experience leaking at first. This does tend to settle but can take several months. Do not become discouraged and please continue with pelvic floor exercises.

We will discharge you soon after your catheter removal when we are happy you are emptying your bladder.

What to expect once you leave hospital

After removal of your catheter it may take some time for you to gain complete control of your bladder. During this time it may be necessary for you to wear a pad in your underwear. We will give you a small supply of pads when we initially remove your catheter and when you are discharged from hospital we will give you the contact details of the Continence Advisory Service, who will arrange further supplies.

If you have not already been given an advice leaflet regarding pelvic floor exercises, please ask one of the ward nurses to get you one. It may take a couple of weeks for you to see any results from these exercises but please continue to do them as they do help.

Contact details

Hopkins Ward Tel. 0118 322 7771 / 7274 (direct line)

Further information

Continence Advisory Service

Tel: 0118 949 5145 (general enquiries) / 0118 949 5146 (clinic enquiries)

Pelvic floor exercises for men

https://prostatecanceruk.org/media/975926/pelvic_floor_exercises-ifm.pdf

Royal Berkshire NHS Foundation Trust

London Road

Reading RG1 5AN

Telephone 0118 322 5111

www.royalberkshire.nhs.uk

This document can be made available in other languages and formats upon request.

Produced by Karen Linger, Urology Team, July 2015

Reviewed by Melanie Martin, July 2019

Review due: July 2021