

Will I be able to go to work?

If possible, please follow your normal daily routine. However, if you prefer not to remain at work, try to be as active as you normally would be at home.

Will I be able to bathe or shower?

The recorder must not get wet so you must not take a bath, shower or go swimming.

Damage to the equipment

Please treat the equipment with care. The catheter is particularly vulnerable and if it catches on a door handle for example, it will be irreversibly damaged. We recommend that you wear a loose top over the recorder and catheter to protect it from snagging.

What if I can't tolerate the catheter?

In this unlikely event, you can remove the catheter yourself. To do this, remove the tape from your face, take a deep breath in and pull it out from your nose. Please dispose of the catheter and return the recorder and diary to the unit. Obviously, if this happens we will gain less information about your condition.

What happens at the end of the test?

At the end of the test the catheter needs to be removed. This is a very simple procedure and the practitioners will teach you how to do this.

The recorder and diary need to be returned promptly, either by hand or by special delivery guaranteed post.

When will I know the result?

The results are sent back to the doctor or consultant who referred you for the tests. You should allow two weeks for the results to reach them.

Glossary of terms

Oesophagus: The oesophagus or gullet is a muscular tube, which extends from the throat to the stomach.

pH: A scale of numbers between 1 and 14 that show the strength of acid or alkali.

Impedance: The detection of either liquid or a mixture of liquid and gas refluxing into the oesophagus from the stomach.

Catheter: Flexible plastic tube.

Reflux: Flow of stomach contents into the oesophagus.

Visit our website www.royalberkshire.nhs.uk

GI Physiology Department

C/o Endoscopy Unit

Royal Berkshire NHS Foundation Trust

Level 2 South Block, London Road


Reading RG1 5AN

GI Physiology Department, January 2018

Review due: January 2020

NHS

Royal Berkshire
NHS Foundation Trust



24 hour oesophageal pH impedance test

Information for patients

GI Physiology Practitioners

0118 322 7725

Why do I need the test?

The test is designed to measure the amount of reflux (both acidic and non-acidic) from your stomach into your oesophagus (gullet) over 24 hours. It also measures anything that has been regurgitated, whether it is liquid or gas and will help you to find out if your symptoms are caused by acid reflux or not.

What to expect before the test

- You should stop taking indigestion medication **one week** before the test so that we can obtain an accurate assessment of what is refluxing into your oesophagus. Full instructions will be sent in your appointment letter as this can vary depending on which drugs you are taking.
- You should continue taking medication for other conditions, e.g. diabetes, epilepsy or blood pressure as normal.
- You should not eat or drink anything in the four hours before the test, apart from sips of water.
- You are advised to wear loose clothing, a separate top and a skirt or trousers, as the catheter is inserted under your top and attached to the machine, which is worn over your shoulder.

What to expect on arrival

The GI Physiology practitioner will explain the procedure to you and ask you to sign a

consent form. This is to ensure that you understand the test and its implications.

The practitioner will also ask you a series of questions about your symptoms and the medication you normally take for them. This will help with the interpretation of the results. Please bring a list of your medications with you.

What to expect during the test

You will be made comfortable on the couch, sitting upright. The practitioner will then pass a very fine measuring catheter (flexible tube) through your nose and into your oesophagus. You will not be asleep or sedated in any way as the procedure should not be painful and you will only feel a slight discomfort. The catheter will remain in place for the duration of the test (about 24 hours) and is attached to a recording machine, worn over your shoulder. You will be asked to complete a diary during the period of the test.

How do I complete the diary sheet?

The practitioner will explain this to you during your visit. Most importantly, we want you to record when you experience your symptoms, e.g. heartburn, regurgitation etc.

Will I be able to feel the catheter?

You will be aware of the catheter in the back of your throat throughout the test but most patients find they become less aware of it with time.

Will I be able to eat and drink?

We especially want you to continue with your normal diet throughout the test as we need to see what happens during a normal day. Please do not chew gum.

The catheter may move very slightly as you eat and it may feel strange but we would like you to persevere as it is important to know what happens after mealtimes.

Will I be able to sleep?

On going to bed, you may remove the recorder from your shoulder and put it (still attached) on your bed or bedside table. Again, it is important that you sleep in your normal position and record any symptoms that disturb you during the night.

Can I drive wearing it?

The medical advice is that you should not drive during the test as you may be distracted. If you are coming by car to the hospital, please make sure you are accompanied by a driver.

What about indigestion medicine?

Please do not take any indigestion medication during the test as all medications for indigestion have the effect of reducing or masking the amount of acid present and therefore will not give an accurate result.